



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 5/00, 5/02	A1	(11) International Publication Number: WO 00/50569 (43) International Publication Date: 31 August 2000 (31.08.00)
(21) International Application Number: PCT/US00/04548 (22) International Filing Date: 23 February 2000 (23.02.00) (30) Priority Data: 60/121,482 24 February 1999 (24.02.99) US (71) Applicant (for all designated States except US): THE WISTAR INSTITUTE OF ANATOMY AND BIOLOGY [US/US]; 3601 Spruce Street, Philadelphia, PA 19104-4268 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SANTOLI, Daniela [US/US]; 933 Wootton Road, Bryn Mawr, PA 19010 (US). ROVERA, Giovanni [US/US]; 933 Wootton Road, Bryn Mawr, PA 19010 (US). CESANO, Alessandra [IT/US]; 226 W. Rittenhouse Square, The Dorchester #413, Philadelphia, PA 19103 (US). (74) Agents: BAK, Mary, E. et al.; Howson and Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHOD OF MODIFYING CYTOTOXIC CELLS AND USES THEREOF		
(57) Abstract		
<p>TALL-104 cells, and other cytotoxic T cell lines, may be modified to increase the cytotoxicity thereof, to enhance growth properties, and/or to provide a preferred phenotype, e.g., expression of cell surface antigens, function, e.g., change in cytokine production profile, by culturing the cells in an effective amount of IL-15, optionally followed by gamma irradiation to halt proliferation.</p>		
Express Mail No. ET033647367US		

ABSTRACT OF THE DISCLOSURE

TALL-104 cells, and other cytotoxic T cell lines, may be modified to increase the cytotoxicity thereof, to enhance growth properties, and/or to provide a preferred phenotype, e.g., expression of cell surface antigens, function, e.g., change in
5 cytokine production profile, by culturing the cells in an effective amount of IL-15, optionally followed by gamma irradiation to halt proliferation.